

Patient's Name _____ D.O.B. _____ Date _____

Referring Physician _____ Phone _____

REFERRING PHYSICIAN'S SIGNATURE _____

Clinical History _____

Diagnosis _____

MRI Check for w/ and w/o contrast

- Abdomen.....
- Adrenal Glands.....
- Brachial Plexus.....
- Brain.....
- Breast.....
- Breast BiopsyLeft Right
- Chest.....
- Coccyx.....
- C-Spine.....
- Enterography.....
- Internal Auditory Canals.....
- Kidneys.....
- Liver.....
- L-S Spine.....
- MRCP.....
- Orbits.....
- Pancreas.....
- Pelvis.....
- Pituitary.....
- Sacro-Iliac Joints.....
- Sacrum.....
- Sinuses.....
- Soft Tissue Neck.....
- TM Joints.....
- T-Spine.....

Extremities

- Ankle..... L R
- Elbow..... L R
- Foot..... L R
- Hip..... L R
- Knee..... L R
- Shoulder.. L R
- Other _____
- Wrist..... L R
- Hand..... L R
- Thigh/Femur.. L R
- Tibia/Fibula.. L R
- Forearm... L R
- Humerus.. L R

MR ANGIOGRAPHY

- Aorta.....
- Carotid.....
- Intracranial.....
- Lower Extremity Left Right
- Renal.....
- Other _____

PET/CT

- Skull Base to Mid-Thigh CPT 78815
- Whole Body CPT 78816
- Brain CPT 78608

CT

- Check for IV contrast**
without with with/without
- Abdomen.....
 - Abdomen & Pelvis.....
 - Brain.....
 - C-Spine.....
 - Chest.....
 - Chest High Res.....
 - Internal Auditory Canals.....
 - L-S Spine.....
 - Low Dose Lung Cancer Screening.....
 - Orbits.....
 - Pelvis.....
 - Sinuses.....
 - Soft Tissue Neck.....
 - T-Spine.....
 - Other_____

CT ANGIOGRAM (CTA)

- CTA Abdomen.....
- CTA Abdomen & Pelvis.....
- CTA Chest.....

Extremities

- Upper:**
- Shoulder.....Left Right
 - Humerus.....Left Right
 - Elbow.....Left Right
 - Forearm.....Left Right
 - Wrist.....Left Right
 - Hand.....Left Right
 - Other _____ Left Right

Lower:

- Hip.....Left Right
- Ankle.....Left Right
- Foot.....Left Right
- Femur.....Left Right
- Knee.....Left Right
- Tibia/Fibula...Left Right
- Other _____ Left Right

BONE DENSITOMETRY

- Bone Density Study
- Vertebral Fracture Assessment

MAMMOGRAPHY/3D MAMMOGRAPHY

- Screening.....LeftRightBilateral
- Diagnostic.....LeftRightBilateral
- Breast Ultrasound if needed

GENERAL ULTRASONOGRAPHY

- Abdomen with Gallbladder
- Aorta Bladder Pelvis
- Doppler Carotid
- Arteries Lower Extremity..Left ..Right ..Bilateral
- Arteries Upper Extremity..Left ..Right ..Bilateral
- Veins Lower Extremity..Left ..Right ..Bilateral
- Veins Upper Extremity..Left ..Right ..Bilateral
- Renal
- Thyroid
- Scrotum w/doppler if necessary
- Other _____

WOMEN'S IMAGING - ULTRASONOGRAPHY

- Breast.....Left Right Bilateral
- OB Biophysical Profile
- OB >14 weeks
- OB Detailed Fetal Anatomy
- OB Transvaginal
- OB Other _____
- Pelvic w/doppler if necessary
- Pelvic/Transvaginal w/doppler if necessary
- Transvaginal w/doppler if necessary
- Sonohysterography
- Other _____

BIOPSIES

- Breast Core Biopsy.....LeftRight
- Breast Cyst Aspiration.....LeftRight
- Stereotactic Breast Biopsy.....LeftRight
- Lymph Node Core Biopsy
- Lymph Node Needle Aspiration
- Soft Tissue Mass Core Biopsy
- Soft Tissue Neck
- Thyroid Fine Needle Aspiration..LeftRight

X-RAY

- Abdomen
- C-Spine
- KUB
- L-S Spine
- Pelvis
- T-Spine
- Other _____
- Chest
- Facial Bones
- Nasal
- Sinuses
- Skull

Extremities

- Ankle.....LeftRight
- Elbow.....LeftRight
- Femur.....LeftRight
- Foot.....LeftRight
- Hand.....LeftRight
- Hip.....LeftRight

- Humerus.....LeftRight
- Knee.....LeftRight
- Radius/Ulna.....LeftRight
- Shoulder.....LeftRight
- Tibia/Fibula.....LeftRight
- Wrist.....LeftRight
- Other.....LeftRight

INSTRUCTIONS FOR EXAMINATION

■Abdomen and/or Pelvic CT Studies

Contact our office to see if oral contrast is required and needs to be picked up in advance.

■CT I.V. Contrast Studies

Nothing to eat or drink for 4 hours prior to exam. Normal medications, that need to be taken within this 4 hour period, should be taken with a small amount of water. Diabetics should have nothing to eat or drink for 3 hours prior to the exam. Patients taking glucophage or any medication containing glu-cophage must discontinue it on the day of the exam, and for 2 days after. The patient must have a stat blood test, to check renal functions, before resuming their medication. All prepara-tions may need to be modified according to diagnosis.

■MRI/MRA

Please come without any metal or jewelry.

MRCP, Routine Abdomen, Liver, Pancreas:

Nothing to eat or drink for 4 hours prior to the exam.

Normal medications that need to be taken within this 4-hour period should be taken with a small amount of water.

ALL OTHER MRI/MRA STUDIES: No preparation. If the patient is going to require sedation, our office must be notified, so that the patient can be given specific instructions prior to taking the sedative.

■PET/CT

If the patient is diabetic, contact our office for instructions.

ON THE DAY BEFORE EXAMINATION:

Remain as calm and quiet as possible. No physical activity. Drink at least 4 glasses of water

DAY OF EXAMINATION:

Remain as calm and quiet as possible. No physical activity. Nothing other than water to eat or drink for 6 hours before the test.

■Mammography

No deodorant or powder on the underarms or breasts the day of the exam.

■Esophagram

Nothing to eat or drink (not even water) for six hours before the test.

■Small Bowel Series

Nothing to eat or drink (not even water) after midnight on the night before the exam, and the morning of the exam. The preparation may need to be modified by the patient's physician.

■Upper Gastrointestinal Series

Nothing to eat or drink (not even water) after midnight the night before the exam and morning of the exam.

■Ultrasound

BLADDER: See "Pelvic" below.

PELVIC: Empty bladder. Drink 6, 8 oz. glasses of water (48 oz.) 45 minutes before the test. Complete drinking in 20 minutes and do not empty bladder.

PREGNANCY-PELVIC (TRANSABDOMINAL) OB: Empty bladder. Drink the following amounts one hour before the test, and do not empty bladder.

6-15 weeks: Drink 5, 8-oz. glasses of water (40 oz.)

16-20 weeks: Drink 2, 8-oz. glasses of water (16 oz.)

Over 20 weeks: No preparation.

PREGNANCY TRANSVAGINAL OB: No preparation.

ABDOMINAL (including gallbladder):

Nothing to eat or drink 6 hours before the exam.

RENAL/RENAL and BLADDER: Nothing to eat or drink 6 hours before exam. 30 minutes before exam, empty bladder, then drink 4, 8 oz. glasses of water (32 oz.) and do not empty bladder.

ALL OTHER ULTRASOUNDS: No preparation necessary.



a **Rezolut** partner

175 Memorial Highway, LL14

New Rochelle, NY 10801

Tel: (914) 633-7700 • Fax: (914) 576-8503

www.newrorad.com

New Rochelle Radiology location:

